arXiv:hep-ph/9909499v1 24 Sep 1999

SUSYGEN3, AN EVENT GENERATOR FOR LINEAR COLLIDERS

N. GHODBANE
Institut de Physique Nucléaire de Lyon
43 Bd du 11 novembre 1918, 69622 Villeurbanne cedex, France

The Monte Carlo program `SUSYGEN`, initially designed for computing distributions and generating events for supersymmetric particle production in e^+e^- collisions, has now been upgraded to study supersymmetric processes at linear colliders by the inclusion of beamstrahlung, beam polarization, spin correlations and complex couplings including CP violating phases. It continues to offer, in the new context, the possibility to study the production and decay of supersymmetric particles within the most general minimal standard supersymmetric model (MSSM), the minimal supergravity model (MSUGRA) or the gauge mediated supersymmetry breaking model (GMSB), with or without assuming R-parity conservation.

1 SUSYGEN2 content and upgrades

Hopefully the linear collider would operate in an environment where a few sparticles would have been discovered and the high statistics will permit a precise study of the soft supersymmetry breaking parameters [1]. One has therefore to move from a *searches* Monte Carlo to a *"precision"* physics Monte Carlo since one would need to measure accurately the above parameters. `SUSYGEN2` [2] is a Monte Carlo event generator for the production and decay of supersymmetric particles and has been initially designed for e^+e^- colliders. It has been extensively used by all four LEP experiments to simulate the expected signals. It includes pair production of charginos and neutralinos, scalar leptons and quarks. It offers also the possibility to study the production of a gravitino plus a neutralino within GMSB models and the production of single gauginos if one assumes R-Parity to be broken All important decay modes of supersymmetric particles relevant to LEP energies have been implemented, including cascades, radiative decays and R-Parity violating decays to standard model particles. The decay of particles is done through the exact matrix elements. The lightest supersymmetric particle (LSP) can either be the neutralino ($\tilde{\chi}_1^0$), the sneutrino ($\tilde{\nu}$) or the gravitino ($\tilde{G}$) in R-Parity conserving models, or there can be no LSP if R-Parity is violated. The initial state radiative corrections take account of p_T/p_L effects in the Structure Function formalism. QED final state radiation is implemented using the `PHOTOS` [4] library. An optimized hadronization interface to `JETSET` [5] is provided, which also takes into account the lifetime of sparticles. Finally, a widely used feature of `SUSYGEN2` is the possibility to do

automatic scans on the parameter space through user friendly ntuples. Recently SUSYGEN2 has been upgraded to SUSYGEN3 [3] in order to adapt to the needs of the next generation of linear colliders, but also in order to extend its potential to supersymmetric particles searches at e^-p colliders (e.g HERA) and hadronic colliders (e.g TEVATRON or LHC). The main new features relevant for linear colliders are the inclusion of beamstrahlung through an interface to CIRCEE [6], the full spin correlation in initial and final states, the inclusion of CP violating phases and the possibility to have an elaborate calculation of the MSUGRA spectrum through an interface to SUSPECT [7].

2 Beam polarization and spin correlations

Since one expects high luminosities for the next generation of linear colliders (e.g. $\sim 500 fb^{-1}$ for the TESLA project), one can use beam polarization to reduce the standard model backgrounds and use the polarization dependence of the cross sections to study specific SUSY parameters. Moreover, as it has been stressed by several authors [8], spin correlations play a major role in the kinematic distributions of final particles. To fulfill these two requirements, the *helicity amplitude method* [9] was used for the calculation of the different feynman amplitudes for production and decay, in order to obtain full spin correlation. Since such an amplitude involves products and contractions of fermionic currents, two basic functions, namely the B and Z functions were defined through:

$$\begin{array}{lcl} B^{\lambda}_{\lambda_1,\lambda_2}(p_1,p_2) & = & \bar{u}_{\lambda_1}(p_1,m_1)P_\lambda u_{\lambda_2}(p_2,m_2), \\ Z^{\lambda\lambda'}_{\lambda_1,\lambda_2,\lambda_3,\lambda_4}(p_1,p_2,p_3,p_4) & = & [\bar{u}_{\lambda_1}(p_1,m_1)\gamma^\mu P_\lambda u_{\lambda_2}(p_2,m_2)] \\ & \times & [\bar{u}_{\lambda_3}(p_3,m_3)\gamma_\mu P_{\lambda'} u_{\lambda_1}(p_4,m_4)]\,, \end{array}$$

where P_λ stands for one of the two chiral projectors P_L or P_R and $u_\lambda(p,m)$ denotes the positive energy spinor solution of the Dirac equation for a particle of helicity λ, four momentum p and mass m. The decomposition of the bispinors $u_\lambda(p,m)$ in terms of the massless helicity eigenstates $\omega_\lambda(k)$ yields analytic and easy to handle expressions for the B and Z functions. The amplitude is then factorized in terms of these basic building blocks; this fact permits compact and transparent coding and speed of calculation. The masses are not neglected in any stage of the calculation. For gaugino productions and decay, we use the *widthless approximation*. Let us illustrate it with, for instance, the calculation of the cross section associated to $e^+e^- \to \tilde{\chi}^0_2\tilde{\chi}^0_1 \to \tilde{\chi}^0_1\tilde{\chi}^0_1 e^+e^-$. The total amplitude associated to a given helicity configuration of the different particles is approximated by the product of the amplitude associated to the production of the two neutralinos ($\tilde{\chi}^0_2\tilde{\chi}^0_1$) with the amplitude corresponding to the decay of

the next to lightest neutralino ($\tilde{\chi}_2^0$). The propagator squared of $\tilde{\chi}_2^0$ is approximated by a factor given by $8\pi^4/(m_{\tilde{\chi}_2^0}\Gamma_{\tilde{\chi}_2^0})$. The total amplitude expression is given by:

$$\mathcal{M} \sim \frac{2\sqrt{2}\pi^2}{m_{\tilde{\chi}_2^0}^{1/2}\Gamma_{\tilde{\chi}_2^0}^{1/2}} \sum_{\lambda_{\tilde{\chi}_2^0}} \mathcal{M}(e^+e^- \to \tilde{\chi}_2^0\tilde{\chi}_1^0) \times \mathcal{M}(\tilde{\chi}_2^0 \to \tilde{\chi}_1^0 e^- e^+)$$

The phase space integration is done through multichannel method [12]. The left side of figure 1, illustrates the effect of spin-correlations in angular distributions of decay electrons for the neutralinos case [?]. Recent studies [10] have shown that τ polarization effects yield valuable information for the MSSM parameters e.g for $\tan\beta$, the nature of the LSP and the mixing angle $\theta_{\tilde{\tau}}$. That is why, we interfaced `SUSYGEN3` to `TAUOLA` [11] which takes into account the τ polarization accurately in the decays. The large effects one expects, for the momenta of the decay distributions are shown on the right side of figure 1.



Figure 1: The figure on the left side shows the $d\sigma/d\cos\theta$ distribution for the final e^- associated to the process $e^+e^- \to \tilde{\chi}_2^0\tilde{\chi}_1^0$ with and without taking into account spin correlations. The figure on the right shows the momentum distribution of e, μ and π, decay products of τ produced in the process $e^+e^- \to \tilde{\tau}_1^+\tilde{\tau}_1^- \to \tau^+\tilde{\chi}_1^0\tau^-\tilde{\chi}_1^0$. The distributions have been plotted for two hypotheses concerning the stau mixing angle $\theta_{\tilde{\tau}}$ ($\tilde{\tau}_L$ and $\tilde{\tau}_R$) and compared to the no spin correlation case

3 Including phases in supersymmetry searches

In the MSSM, there are new potential sources of CP non conservation[13]. Complex CP violating phases can arise from several parameters present in the MSSM Lagrangian: the higgs mixing mass parameter μ, the gauginos masses M_i, the trilinear couplings A_i. Experimental constraints on these CP violating phases come from the electric dipole moment of the electron and the neutron. Since all the couplings [14] and the different mass parameters μ, M_1, and the trilinear couplings A_τ, A_t and A_b have been assumed to be complex by default, the introduction of phases in the gaugino and sfermion sector for masses as well for cross sections has been straightforward in `SUSYGEN3`. Figure 2 shows the chargino pair production cross section variations in terms of ϕ_μ, the phase associated to the μ parameter, for several values of the sneutrino mass $m_{\tilde{\nu}_e}$ and for a value of $\tan\beta$ equal to 1. One sees that there is a local minimum between the two extreme values ($\Phi_\mu = 0, 180$ degrees), by default examined at LEP searches, but also that it is not so deep as to raise doubts on the exhaustiveness of "phaseless" searches. On the contrary at the linear collider, one cannot neglect this strong dependence of the cross section from phases.



Figure 2: $e^+e^- \to \tilde{\chi}_1^+\tilde{\chi}_1^-$ cross section evolution in terms of ϕ_μ, the phase associated to the higgsino mixing mass parameter μ. We also assume different hypotheses concerning the sneutrino $\tilde{\nu_e}$ mass

4 Future projects

Native C++ event generators are needed to optimize the couplings with linear collider simulation based on `GEANT 4` and analysis tools. This will enable to

share code and reuse classes between different generators. The helicity amplitude method and the two basic functions, namely the B and the Z functions, used to upgrade `SUSYGEN` seem to be optimally suited for object oriented implementation. Moreover, *Multichannel integration* looks well adapted for object treatment.

1. See e.g. H.E. Haber and G.L. Kane, Phys. Rept. **117** 75 (1985),
2. S. Katsanevas and P. Morawitz,*Comput. Phys. Commun.* **112**, 227 (1998),
3. N. Ghodbane, S. Katsanevas, P. Morawitz and E. Perez, `SUSYGEN3` `http://lyoinfo.in2p3.fr/susygen/susygen3.html`
4. E. Barberio, Z. Was, *Comput. Phys. Commun.* **79**, 291 (1994),
5. T. Sjöstrand, *Comput. Phys. Commun.* **82**, 74 (1994),
6. T. Öhl, *Comput. Phys. Commun.* **101**, 1997 (269),
7. A. Djouadi, J.L. Kneur and G. Moultaka, Note GDR–S–17,
8. G. Moortgat-Pick *et al*, hep-ph/9708481, S. Y. Choi *et al*, hep-ph/9806279, and V. Lafage *et al*,hep-ph/9810504, see also D. Perret-Gallix: Automatic amplitude calculation and event generation for linear colliders: `GRACE` and `COMPHEP`, these proceedings,
9. M. Martinez, R. Miquel and C. Mana, *Helicity amplitudes calculation.* Proceedings on the Workshop on QED Structure Functions, Ann Arbor, Michigan, 22-25 May 1989, p. 24,
10. M. Nojiri, hep-ph/9511338, A. Bartl *et al*, Z. Phys. **C73**, 469 (1997)
11. S. Jadach *et al Comput. Phys. Commun.* **76**, 1993 (361),
12. R.Kleiss and R.Pittau, Comput. Phys. Commun. 83 141 (1994),
13. see e.g. T. Ibrahim, P. Nath, Phys.Rev. D58 111301 (1998),T. Falk and K.A. Olive, Phys. Lett. **B439**, 71 (1998), M. Brhlik, G.J. Good and G.L. Kane, Phys. Rev. **D59**, 115004 (1999), S. Pokorski, J. Rosiek, C.A. Savoy, hep-ph/9906206 v3,
14. J. Rosiek, *Phys. Rev.* D **41**, 3464 (1990) and hep-ph/9511250 (erratum).


σ(e+e-→ τ̃1τ̃1) (pb)
0.7
0.6
0.5
0.4
0.3
0.2
0.1
0
e- beam polarisation -1
e- beam not polarised
e- beam polarised +1
0
10
20
30
40
50
60
70
80
90
100
fraction of e+ beam polarized +1 (in percent)


Cross section evolution in terms of the μ phase
σ(e+e-→ χ̃1+χ̃1-) (pb)
M2=80 GeV
tanβ=1.5
M(ν̃)=80 GeV
μ= 50
μ= 150
μ= 200
Φμ
0
1
2
3
4
5
6
7
8
0
20
40
60
80
100
120
140
160
180


400
350
300
250
200
150
100
50
0
0
0.02
0.04
0.06
0.08
0.1
0.12
0.14
0.16
0.18
0.2
E_π/E_{beam}
$\tau \rightarrow \nu_\tau \pi$
$\tilde{\tau}_1=\tilde{\tau}_L$
$\tilde{\tau}_1=\tilde{\tau}_R$
No H.E.


P_τ
1
0.8
0.6
0.4
0.2
0
-0.2
-0.4
-0.6
-0.8
-1
0
50
Θ_τ
tanβ=3
tanβ=15
tanβ=30
0
20
40
tanβ
Θ_τ=0
Θ_τ=π/4
Θ_τ=π/2